

# PANORAMA

Thematic portfolio



## Controlling bovine tuberculosis: a One Health challenge



# PERSPECTIVES

# DOSSIER

# AROUND THE WORLD

*Forty-four percent of countries reported bovine tuberculosis (bTB) via the OIE World Animal Health Information System (WAHIS) between January 2017 and June 2018. Only a quarter of the affected countries were applying all the relevant control measures. Improved surveillance and accurate reporting by a country's Veterinary Services contributes to the prevention and control of bTB at the animal source.*

**From January 2017 to June 2018**, of the 188 countries and territories reporting their bTB situation to the OIE, 82 countries (44%) were affected, which demonstrates a widespread distribution of the disease (Fig. 1).




Fig. 1 – Global distribution of bovine tuberculosis in 2017 and first semester of 2018

Among the 82 affected countries, 29 (35.4%) countries reported the presence of bTB in both livestock and wildlife. Two (2.4%) countries reported bTB present only in wildlife, compared to 51 (62.2%) which indicated that only livestock were affected. Moreover, among these 82 affected countries, 66 (80.5%) provided quantitative data for outbreaks via WAHIS, demonstrating relatively good reporting of the global situation of this disease. The persistence of the infection in wildlife poses challenges for disease control in some countries [1], due to the potentially significant impact of wildlife as reservoirs and spillover hosts.

## What are we doing to fight bovine tuberculosis?

The implementation of relevant measures is crucial for preventing and controlling bTB at its animal source to avoid its transmission between animals and to humans.





© OIE

An analysis of the implementation of prevention and control measures for bTB by country, using WAHIS data, showed that, among the affected countries, 23% had implemented all the relevant control measures. These were: active surveillance, complete or partial stamping-out, and movement control. The majority (62%) of affected countries had implemented some of the relevant measures. However, 3% of them had not applied any of these measures, and therefore require strengthened control efforts.

Among the countries where the disease was reported as absent, the majority (82%) recorded implementing at least one of the relevant prevention measures, namely surveillance and/or border control.

The above data demonstrate a high level of vigilance for bTB by affected and non-affected countries. As outlined in the *Roadmap for Zoonotic Tuberculosis* [2], the collection and reporting of more complete and accurate data is considered one of the priority areas for controlling zoonotic tuberculosis caused by *Mycobacterium bovis*. For this reason, countries are encouraged to continue to maintain and improve their level of bTB surveillance and reporting.

<http://dx.doi.org/10.20506/bull.2019.1.2912>

## DOSSIER

# Bovine tuberculosis: global distribution and implementation of prevention and control measures according to WAHIS data

### KEYWORDS

#bovine tuberculosis, #control measure, #global distribution, #map, #OIE World Animal Health Information System (OIE-WAHIS), #prevention measure, #World Organisation for Animal Health (OIE).

### AUTHORS

Kiyokazu Murai<sup>(1)\*</sup>, Paolo Tizzani<sup>(2)</sup>, Lina Awada<sup>(2)</sup>, Lina Mur<sup>(2)</sup>, Neo J. Mapitse<sup>(3)</sup> & Paula Caceres<sup>(4)</sup>


(1) Chargé de mission, World Animal Health Information and Analysis Department, World Organisation for Animal Health (OIE).

(2) Veterinary Epidemiologist, World Animal Health Information and Analysis Department, World Organisation for Animal Health (OIE).

(3) Head, Status Department, World Organisation for Animal Health (OIE).

(4) Head, World Animal Health Information and Analysis Department, World Organisation for Animal Health (OIE).

\* Corresponding author: [k.murai@oie.int](mailto:k.murai@oie.int)



© P. Bastiaensen

## REFERENCES

1. Palmer M.V. (2013). – *Mycobacterium bovis*: characteristics of wildlife reservoir hosts. *Transbound. Emerg. Dis.*, **60** (Suppl. 1), S1–13.  
<https://doi.org/10.1111/tbed.12115>.
2. World Health Organization (WHO), World Organisation for Animal Health (OIE) & Food and Agriculture Organization of the United Nations (FAO) (2017). – [Roadmap for Zoonotic Tuberculosis](#).

**The OIE** is an international organisation created in 1924 with a mandate from its 182 Member Countries to improve animal health and welfare. Its activities are permanently supported by 301 centres of scientific expertise and 12 regional offices with a presence on every continent.



Follow the OIE: [www.oie.int](http://www.oie.int)



@OIEAnimalHealth



World Organisation for Animal Health - OIE



OIEVideo



World Organisation for Animal Health



World Organisation for Animal Health (OIE)



Digital version: [www.oiebulletin.com](http://www.oiebulletin.com)



**WORLD ORGANISATION FOR ANIMAL HEALTH**

*Protecting animals, preserving our future*

12, rue de Prony - 75017 Paris, France

Tel.: +33 (0)1 44 15 18 88 - Fax: +33 (0)1 42 67 09 87 - [oie@oie.int](mailto:oie@oie.int)